

This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

The following resources related to this article are available online at www.sciencemag.org (this information is current as of July 19, 2011):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/298/5600/1987.full.html>

Supporting Online Material can be found at:

<http://www.sciencemag.org/content/suppl/2002/12/04/298.5600.1987.DC1.html>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/content/298/5600/1987.full.html#related>

This article **cites 31 articles**, 5 of which can be accessed free:

<http://www.sciencemag.org/content/298/5600/1987.full.html#ref-list-1>

This article has been **cited by** 159 article(s) on the ISI Web of Science

This article has been **cited by** 19 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/content/298/5600/1987.full.html#related-urls>

This article appears in the following **subject collections**:

Ecology

<http://www.sciencemag.org/cgi/collection/ecology>

5. R. Cahn, M. Winter, *Indiana* **13**, 30 (1993).
6. G. Whittaker, thesis, Yale University (1980).
7. W. F. Rust, R. J. Sharer, *Science* **242**, 102 (1988).
8. M. Coe, *America's First Civilization* (American Heritage, New York, 1968).
9. R. González Lauck, in *Olmec Art of Ancient Mexico*, E. P. Benson, B. de la Fuente, Eds. (National Gallery of Art, Washington, DC, 1996), pp. 73–81.
10. K. O. Pope et al., *Science* **292**, 1370 (2001).
11. P. Drucker, *La Venta, Tabasco: A Study of Olmec Ceramics and Art*, Smithsonian Institution Bureau of American Ethnology Bulletin 153 (1952).
12. J. Quirarte, in *The Origins of Maya Civilization*, R. E. W. Adams, Ed. (Univ. New Mexico Press, Albuquerque, NM, 1977), pp. 249–283.
13. J. F. Scott, *The Danzantes of Monte Albán*, Studies in Pre-Columbian Art and Archaeology 19 (Dumbarton Oaks, Washington, DC, 1978).
14. E. Benson, B. de la Fuente, Eds., *Olmec Art of Ancient Mexico* (National Gallery of Art, Washington, DC, 1996), p. 213.
15. J. Clark, M. Pye, Eds., in *Olmec Art and Archaeology in Mesoamerica* (National Gallery of Art, Washington, DC, 2000), p. 225.
16. T. A. Lee Jr., *The Artifacts of Chiapa de Corzo, Chiapas, Mexico*. Papers of the New World Archaeological Foundation 26 (Brigham Young Univ., Provo, UT, 1969).
17. P. Agrinier, *The Archaeological Burials at Chiapa de Corzo and Their Furniture*, Papers of the New World Archaeological Foundation 12 (Brigham Young Univ., Provo, UT, 1964).
18. G. W. Lowe, in *The Archaeological Burials at Chiapa de Corzo and Their Furniture*, P. Agrinier, Papers of the New World Archaeological Foundation 12 (Brigham Young Univ., Provo, UT, 1964).
19. M. Coe, personal communication.
20. M. Macri, personal communication.
21. S. Méluzin, *Further Investigations of the Tuxtla Script: An Inscribed Mask and the La Mojarra Stela 1*, Papers of the New World Archaeological Foundation 65 (Brigham Young University, Provo, UT, 1995).
22. R. Berger, J. A. Graham, R. F. Heizer, *Contrib. Univ. Calif. Archaeolog. Res. Facil.* **3**, 1 (1967).
23. L. Campbell, T. Kaufman, *Am. Antiq.* **41**, 80 (1976).
24. L. Schele, M. E. Miller, *The Blood of Kings. Dynasty and Ritual in Maya Art* (Kimball Art Museum, Fort Worth, TX, 1986).
25. V. G. Smith, *Izapa Relief Carving: Form, Content, Rules for Design, and Role in Mesoamerican Art History and Archaeology*, Studies in Pre-Columbian Art and Archaeology 27 (Dumbarton Oaks, Washington, DC, 1984).
26. A. Demarest, *Mesoamérica* **5** (7), 53 (1984).
27. J. A. Graham, L. Benson, *Arqueología* **3**, 77 (1990).
28. J. Urcid Serrano, *Zapotec Hieroglyphic Writing*, Studies in Pre-Columbian Art and Archaeology 34 (Dumbarton Oaks, Washington, DC, 2001).
29. F. Winfield Capitaine, *La Estela 1 de La Mojarra, Veracruz, México* (Center for Maya Research, Washington, DC, 1988).
30. We thank V. Fields, E. Boot, J. Justeson, T. Kaufman, J. Urcid, C. Brown, N. Hopkins, and A. Moreno for their help in analyzing the early writing. A. Moreno drew the San Andrés artifacts, K. Womble redrew the Chiapa de Corzo glyph, and T. Carmona did the photography. J. Urcid provided the San José Mogote illustration for Fig. 1. A. Perrett and E. de Armas helped assemble the illustrations. R. González Lauck and J. Clark supported the excavations, which were funded by the National Science Foundation (SBR-9617389) and the Foundation for the Advancement of Mesoamerican Studies, Inc.

16 September 2002; accepted 25 October 2002

Grassland Responses to Global Environmental Changes Suppressed by Elevated CO₂

M. Rebecca Shaw,^{1*} Erika S. Zavaleta,^{1,2‡} Nona R. Chiariello,³ Elsa E. Cleland,^{1,2} Harold A. Mooney,² Christopher B. Field¹

Simulated global changes, including warming, increased precipitation, and nitrogen deposition, alone and in concert, increased net primary production (NPP) in the third year of ecosystem-scale manipulations in a California annual grassland. Elevated carbon dioxide also increased NPP, but only as a single-factor treatment. Across all multifactor manipulations, elevated carbon dioxide suppressed root allocation, decreasing the positive effects of increased temperature, precipitation, and nitrogen deposition on NPP. The NPP responses to interacting global changes differed greatly from simple combinations of single-factor responses. These findings indicate the importance of a multifactor experimental approach to understanding ecosystem responses to global change.

Human actions are affecting many aspects of the Earth system. The composition of the atmosphere, the climate, the abundance of invasive species, and the area of managed landscapes have all undergone important changes in the past century. These changes are likely to be even greater in this century (1). In almost any setting, realistic global change is decidedly multifactorial. Warming, increased precipitation, increased deposition

of N-containing compounds, and increased atmospheric CO₂ are all likely consequences of CO₂ emissions from fossil fuel combustion and land use change (2). In the past century, atmospheric CO₂ concentration has increased globally by more than 30% (2). Temperature, precipitation, and deposition of biologically available N have increased for large regions (3–5). Further increases in the future are almost certain.

Most of the experimental research on ecosystem responses to global change has addressed responses to single global changes, with relatively few studies exploring responses to two or more interacting treatments (6–8). Experimental manipulations of both temperature and CO₂ concentration are rare at the ecosystem scale (9, 10), even though elevated CO₂ is a primary driver of climate change (2).

Several modeling studies have addressed ecosystem responses to multifactor global changes (11, 12), but the theoretical

foundation for predicting ecosystem responses to simultaneous changes in multiple factors is incomplete. For some processes (such as photosynthesis), well-tested mechanistic models support the simulation and interpretation of multifactor responses (13). For many other processes, however, including biomass allocation, the timing of seasonal activity, and species replacements, the empirical data are too sparse to support credible models or allow comprehensive hypothesis tests.

Both empirical and modeling studies highlight potential contrasts in responses to single global changes and multiple, interacting global changes. Stimulation of plant growth by elevated CO₂, for example, may be strongest when water is limiting (14), when nutrients are abundant (15), or when plant species diversity is high (6). Simulated ecosystem responses to future global changes depend strongly on such interactions. In many settings, simulated warming increases decomposition more than net primary production (NPP), leading to a loss of carbon (16). In others, elevated CO₂ and N deposition tend to increase NPP more than decomposition, leading to carbon storage. In some simulations, the responses of carbon storage to the three factors nearly cancel one another out. In others, changes combine in nonadditive ways, with examples of both suppression and amplification (11).

One of the keys to understanding the long-term impacts of multiple global changes on ecosystem function will be experiments on model ecosystems that are amenable to factorial manipulations and respond rapidly. Annual grassland, with a high diversity of small short-lived plants, is an attractive model system for global change experiments. An area of less than 1 m² is sufficient for a meaningful global-

¹Department of Global Ecology, Carnegie Institution of Washington, 260 Panama Street, Stanford, CA 94305, USA. ²Department of Biological Sciences, Stanford University, Stanford, CA 94305, USA. ³Jasper Ridge Biological Preserve, Stanford University, Stanford, CA 94305, USA.

*To whom correspondence should be addressed. E-mail: shaw@globalecology.stanford.edu

†Present address: The Nature Conservancy of California, 201 Mission Street, 4th floor, San Francisco, CA 94105–1832, USA.

‡Present address: The Nature Conservancy of California and Department of Integrative Biology, University of California, Berkeley, CA 94720, USA.





Fig. 1. (A) Schematic drawing of the study plots, side view (left) and top view (right). The plot is 2 m in diameter. (B) Photograph of a study plot.

Fig. 2. NPP. Open bars represent all treatment combinations with ambient CO₂. Gray bars represent all treatment combinations with elevated CO₂. Values are means ± 1 SE, N = 64 plots.



change manipulation of a complete ecosystem with thousands of plants and a full suite of plant, animal, microbial, and soil processes. The annual life cycle of the dominant plants allows relatively brief experiments to span several complete generations (17).

We investigated the impacts of individual and multiple simultaneous global changes on NPP of a moderately fertile California annual grassland (18), using the Jasper Ridge Global Change Experiment (JRGCE). The JRGCE involved four global change factors at two levels: CO₂ [ambient and 680 parts per million (ppm)], temperature (ambient and ambient plus 80 W m⁻² of thermal radiation), precipitation (ambient and 50% above ambient plus 3-week growing season elongation), and N deposition (ambient and ambient plus 7 g of N m⁻² year⁻¹) in a complete factorial design. The JRGCE, initiated in 1998, includes

eight replicates of each of the 16 treatments. These are organized as a split-plot design with 32 circular plots, each divided into four 0.78-m² quadrants, separated by solid partitions below ground and mesh partitions above ground (Fig. 1) (19). Manipulations began at the start of the 1998–1999 growing season. Each year, we estimated NPP as peak biomass (above and below ground) in late spring (20).

Over the first 2 years of the manipulations and across all of the treatment combinations, elevated CO₂ had no significant effect on NPP (21) (Fig. 2). In the third year, 2001, the mean NPP for all treatment combinations with elevated CO₂ was 988 ± 52 g m⁻² versus 1089 ± 54 g m⁻² for all treatment combinations with ambient CO₂ (P = 0.081). The absence of significant biomass increases in response to elevated CO₂ is consistent with several prior studies at the ecosystem scale (8, 9, 14, 22–24) but contrasts sharply with

the results of most greenhouse experiments and many ecosystem studies, where elevated CO₂ leads to increased aboveground growth (25–28).

In the third year of manipulations of the JRGCE, elevated CO₂ stimulated aboveground biomass in the treatment in which all of the other factors were at ambient levels. Aboveground biomass in the ambient CO₂ treatment was 469 ± 27 g m⁻², and adding CO₂ increased this to 622 ± 24 g m⁻² (P = 0.003, table S1) (29). This comparison parallels results of many other single-factor CO₂ experiments. The 32.6% increase in aboveground biomass is comparable to the 25% increase observed in North Carolina pine plantations (25) and the 20 to 43% increase in crop plants at the Arizona Free-Air CO₂ Enrichment (FACE) site (15, 26).

Each of the treatments involving increased temperature, N deposition, or precipitation (alone or in combination) tended to increase aboveground biomass and NPP, but elevated CO₂ consistently dampened these increases (Fig. 3) (29). The three-factor combination of increased temperature, precipitation, and N deposition produced the largest stimulation of NPP (84%, P = 0.021), but the addition of CO₂ reduced this to 40% (P = 0.028). The suppressive effect of elevated CO₂ was even clearer for below ground biomass, where the average effect across all treatments was a decrease of 22% (P = 0.005) (21).

The NPP response of this ecosystem to multiple global changes was not a simple combination of responses to individual global change factors. Some treatment combinations (such as warming and precipitation) led to intermediate responses, with NPP near the average of the single-factor responses (Fig. 3). Others (such as warming and N deposi-

Fig. 3. Percent changes in NPP for each treatment, relative to corresponding reference. For each treatment combination, the reference is defined as all ambient CO₂ plots in which the variable(s) that defines each treatment combination are also at ambient. amb, ambient; C, elevated CO₂; N, N deposition; T, increased temperature; P, increased precipitation. For treatments T, N, and P, *n* = 24 plots. For treatments TP, NP, and TN, *n* = 12. For treatments C and TNP, *n* = 6. For example, the open bar for the increased temperature pair (T) is calculated using all treatments with increased temperature but not elevated CO₂ (*n* = 24). The gray bar in the same pair is calculated using all treatments with increased temperature and elevated CO₂ (*n* = 24).





Fig. 4. The effect of elevated CO₂ on NPP in relation to NPP under ambient CO₂ for the eight treatment pairs with and without elevated CO₂ (30) for each of the 3 years of treatment. Each point is the mean for all replicates (*n* = 8) in one treatment pair in 1 year.



tion) were approximately additive (Fig. 3). Across the seven treatment combinations involving elevated CO₂ and at least one other factor (30), the addition of elevated CO₂ decreased NPP (*P* = 0.048) (31), which is the opposite of the effect of CO₂ as a single factor (Fig. 3).

Thousands of published papers describe plant or ecosystem responses to elevated CO₂ (32). Generally, stimulatory responses in individual plants (27) often persist at the ecosystem scale, in the form of increased aboveground growth. Relatively few studies address ecosystem NPP responses (above- and belowground). Those that do include examples of substantial NPP increases as well as modest responses and no change (33). Only a few studies address the effects of elevated CO₂ in combination with other global changes at the ecosystem

scale. Oechel *et al.* (9) found that warming could extend the time over which elevated CO₂ stimulated NPP in arctic tundra. Oren and colleagues (8) observed a parallel result with N addition in a pine plantation. Reich *et al.* (6) reported increased sensitivity of NPP to elevated CO₂ with added N, but only when species diversity was high. Several studies from agricultural ecosystems also demonstrate that enhanced N availability can lead to larger increases in NPP in response to elevated CO₂ (34). In all these studies on N-limited ecosystems, the sensitivity of NPP to elevated CO₂ increased with the addition of N or, in the tundra example, with a treatment that indirectly increased N availability.

The JRGCE results suggest a fundamentally different kind of ecosystem response. Previous studies indicate that enhanced

availability of a limiting resource such as N increases an ecosystem's potential to take advantage of elevated CO₂. Our results are more consistent with the idea that, under some circumstances, elevated CO₂ can constrain potential NPP. Several aspects of the JRGCE might account for this contrast. To our knowledge, this is the only ecosystem-scale experiment to date in which the treatments have been applied through several generations of the dominant plants. It is one of very few experiments to explore interactions between warming and elevated CO₂ at the ecosystem scale. It is also one of few ecosystem experiments to simulate N deposition with nitrate (the primary form delivered from the atmosphere in most locations) as opposed to a mixture of nitrate and ammonium.

Several mechanisms could contribute to the suppressive effects of elevated CO₂ reported here. First, it is possible that the responses depend on specific features of the treatments. Simulating N deposition with nitrate may shift the plants from ammonium to nitrate nutrition, which may make them less sensitive to elevated CO₂ than those grown on ammonium (35). The response of grassland NPP to elevated CO₂ is often greatest when water is limiting (36). CO₂-driven alleviation of water stress through decreases in transpiration may nullify any stimulatory effect of added precipitation. Although both of these mechanisms potentially account for a decrease in CO₂ sensitivity, neither is likely to result in a suppressive effect of elevated CO₂.

Second, it is possible that the suppressive effect of CO₂ is a consequence of changes in plant species richness or other aspects of plant community composition. The structure of the plant community was dynamic in the JRGCE. However, the changes in plant species richness and abundance were similar in the single- and multifactor CO₂ treatments (37), arguing against a major role for community composition in explaining the suppressive effect of elevated CO₂ on NPP in multifactor treatments only.

A third possibility is that the suppressive effect of elevated CO₂ is a consequence of limitation by some soil nutrient. This could result from gradual immobilization of a limiting nutrient by soil microorganisms under elevated CO₂, reflecting increased demand by microbes more richly supplied with carbon lost from roots (38–40). Several indicators of soil metabolic activity were enhanced under elevated CO₂ (41, 42). Nutrient limitation under elevated CO₂ could also be a consequence of decreased root allocation, which tends to limit the capacity of plants to forage for nutrients. The effects of decreased root alloca-

REPORTS

tion should be most severe in settings where NPP is high, because NPP drives plant resource demand. In most settings, plants respond to nutrient limitation with an increase in root allocation (43). In the results presented here, however, root allocation consistently decreased in response to elevated CO₂, a pattern that may or may not effectively balance nutrient supply and demand (44).

Previous studies at Jasper Ridge grasslands (45) and many other ecosystems documenting the potential for limitation by phosphorus make it an interesting candidate as a limiting nutrient. It is also possible that N becomes limiting when NPP is increased and CO₂ is elevated. N limitation in the treatments with N deposition, however, would require that the additional N be only sparingly available, as a result of either its form or the timing of its application.

Some of the strongest evidence for the nutrient limitation mechanism comes from the multiyear response. Over all 3 years of manipulations in the JRGCE and all treatment combinations, elevated CO₂ tended to be stimulatory when NPP was low and suppressive when NPP was high (Fig. 4). In the eight pairs of treatment combinations with ambient and elevated CO₂, NPP under ambient CO₂ explained 53% of the effect of adding CO₂, with consistent suppressive effects at levels of NPP greater than 800 g m⁻². This result, closely paralleling that for several ecosystems studied with single-factor manipulations (24), provides further support for the hypothesis that negative effects of elevated CO₂ reflect constraints from a resource that becomes increasingly limiting as demand for it increases.

We do not yet know the contribution of each of these mechanisms to the overall response, but probably all contribute. Experiments to test the involvement of each are now underway.

This report of suppressive effects of elevated CO₂ has a strong message for global change research. Ecosystem responses to realistic combinations of global changes are not necessarily simple combinations of the responses to the individual factors. Accurate predictions of ecosystem responses to suites of global changes depend on successful integration across a range of processes and time scales. Multifactor experiments on ecosystems that are easy to manipulate can provide a rich source of examples as well as test beds for exploring hypotheses with the potential to explain the responses of a wide range of ecosystems. Future experiments should develop theoretical and empirical frameworks for integrating information from these model ecosystems with information from less-easily studied ecosystems that play important roles in the carbon cycle.

References and Notes

- P. M. Vitousek, H. A. Mooney, J. Lubchenco, J. M. Melillo, *Science* **277**, 494 (1997).
- International Panel on Climate Change, *Working Group 1 Third Assessment Report* (Cambridge Univ. Press, Cambridge, MA, 2001).
- T. R. Karl, in *The Regional Impacts of Climate Change: An Assessment of Vulnerability*, R. T. Watson, M. C. Zinyowera, R. H. Moss, D. J. Dokken, Eds. (Cambridge MA, Univ. Press, Cambridge, 1998), pp. 412–425.
- D. R. Easterling et al., *J. Geophys. Res.* **105**, 20101 (2000).
- P. M. Vitousek et al., *Ecol. Appl.* **7**, 737 (1997).
- P. B. Reich et al., *Nature* **410**, 809 (2001).
- E. H. DeLucia et al., *Science* **284**, 1177 (1999).
- R. Oren et al., *Nature* **411**, 469 (2001).
- W. C. Oechel et al., *Nature* **371**, 500 (1994).
- N. T. Edwards, R. J. Norby, *Plant Soil* **206**, 85 (1998).
- A. D. McGuire et al., *Global Biogeochem. Cycles* **15**, 183 (2001).
- D. Schimel et al., *Science* **287**, 2004 (2000).
- G. D. Farquhar, S. von Caemmerer, J. A. Berry, *Planta* **149**, 78 (1980).
- C. E. Owensby, J. M. Ham, A. K. Knapp, L. M. Auen, *Global Change Biol.* **5**, 497 (1999).
- B. A. Kimball, J. R. Mauney, *Agron. J.* **85**, 706 (1993).
- P. M. Cox, R. A. Betts, C. D. Jones, S. A. Spall, I. J. Totterdell, *Nature* **408**, 184 (2000).
- C. B. Field, N. R. Chiariello, in *Earth Systems: Processes and Issues*, W. G. Ernst, Ed. (Cambridge Univ. Press, Stanford, CA, 2000), pp. 297–314.
- Research was conducted at Stanford University's Jasper Ridge Biological Preserve in central California (37°24'N, 122°14'W, elevation 150 m) between October 1998 and June 2001. The grassland is dominated by annual grasses (*Avena barbata* and *Bromus hordeaceus*) and forbs (*Geranium dissectum* and *Erodium botrys*). Typical of a Mediterranean climate, the growing season (October to May) is cool and wet, and the summer (June through September) is hot and dry.
- Control, warming, elevated CO₂, and warming plus elevated CO₂ treatments were each applied to eight plots. Infrared heat lamps were suspended over the centers of the warming plots, heating the plant canopy by 0.8° to 1°C. Atmospheric CO₂ concentrations were elevated to ~680 ppm with a ring of free-air emitters surrounding each plot and delivering pure CO₂ at the canopy level. Control, N deposition, increased precipitation, and N plus increased precipitation treatments were randomly assigned to the four quadrants in every plot. We applied N as Ca(NO₃)₂ at the rate of 7 g of N m⁻² year⁻¹, with an initial application of 2 g of N m⁻² at the first rains and a second application of 5 g of N m⁻² applied as slow-release fertilizer in the first week of January each year. We augmented ambient precipitation events by 50% with drip irrigation and overhead sprinklers and added an additional two rain events in the spring to extend the rainy season by approximately 3 weeks. After 1 year of preliminary measurements, we began treatments in November 1998. Each of the 16 possible combinations of CO₂, temperature, N deposition, and precipitation was replicated 8 times.
- We measured aboveground biomass on two dates, 1 month apart, by harvesting all live and senesced vegetation and litter within a designated harvest area. We report the greater of the two harvests for each treatment. Three 22-mm-diameter soil cores made to a depth of 15 cm were taken within the aboveground harvest area, 2 days after the first aboveground harvest. Soils were wet-sieved through 0.5-mm mesh. Live roots were separated from dead roots. We did not account for additional NPP in root turnover, which, on the basis of other studies at this site, may increase annual root production by 44 to 54% (46).
- We analyzed NPP, aboveground biomass, and belowground biomass using a split-plot analysis of variance with two levels each of CO₂, warming, precipitation, and temperature. The full model is described in (37). The data were logarithmically transformed.
- C. Körner, M. Diemer, B. Schappi, P. Niklaus, J. Arnone, *Int. J. Ecol.* **18**, 165 (1997).
- B. G. Drake, G. Peresta, E. Beugeling, R. Matamala, in *Carbon Dioxide and Terrestrial Ecosystems*, G. W. Koch, H. A. Mooney, Eds. (Academic Press, San Diego, CA, 1996), pp. 197–214.
- G. W. Koch, H. A. Mooney, in *Carbon Dioxide and Terrestrial Ecosystems*, G. W. Koch, H. A. Mooney, Eds. (Academic Press, San Diego, CA, 1996) pp. 415–429.
- E. H. DeLucia et al., *Science* **284**, 1177 (1999).
- B. A. Kimball et al., *Global Change Biol.* **1**, 429 (1995).
- B. A. Kimball, *Agron. J.* **75**, 779 (1983).
- H. Poorter, C. Roumet, B. D. Campbell, in *Carbon Dioxide, Populations, and Communities*, C. Körner, F. A. Bazzaz, Eds. (Academic Press, San Diego, CA, 1996), pp. 375–412.
- A Student's *t* test paired by block was performed on logarithmically transformed data. See supporting online material available on Science Online.
- The 16 experimental conditions can be viewed as 8 pairs in which the members of a pair are identical except that one set of replicates experiences ambient CO₂ and one set experiences elevated CO₂. Seven of these eight pairs have elevated levels of one or more of the following factors: warming, precipitation, or N inputs (see supporting online material).
- A Student's *t* test paired by block was performed to determine the effect of elevated CO₂ on NPP under all treatment combinations, except control and elevated CO₂ alone. This was a posthoc test of the hypothesis that, in the presence of at least one other global change factor, the effect of elevated CO₂ was to decrease NPP. The data were logarithmically transformed.
- C. Körner, *Ecol. Appl.* **10**, 1590 (2000).
- H. A. Mooney et al., in *The Terrestrial Biosphere and Global Change: Implications for Natural and Managed Ecosystems*, B. H. Walker, W. L. Steffen, J. Canadell, J. S. I. Ingram, Eds. (Cambridge Univ. Press, Cambridge, MA, 1999), pp. 141–189.
- K. E. Idso, S. B. Idso, *Agric. For. Meteorol.* **69**, 153 (1994).
- A. J. Bloom, D. R. Smart, D. T. Nguyen, P. S. Searles, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 1730 (2002).
- M. Volk, P. A. Niklaus, C. Körner, *Oecologia* **125**, 380 (2000).
- E. S. Zavaleta, thesis, Stanford University (2001).
- S. Hu, I. F. S. Chapin, M. K. Firestone, C. B. Field, N. R. Chiariello, *Nature* **409**, 188 (2001).
- Z. G. Cardon et al., *Soil Biol. Biochem.* **33**, 365 (2001).
- S. Diaz, *Plant Soil* **187**, 309 (1996).
- Y. Luo, R. B. Jackson, C. B. Field, and H. A. Mooney, *Oecologia* **108**, 130 (1996).
- B. A. Hungate, C. H. Jaeger III, G. Gamara, F. S. Chapin III, C. B. Field, *Oecologia* **124**, 589 (2000).
- F. S. Chapin III, A. J. Bloom, C. B. Field, R. H. Waring, *Bioscience* **37**, 49 (1987).
- Y. Luo, C. B. Field, H. A. Mooney, *Plant Cell Environ.* **17**, 1195 (1994).
- R. T. Koide, L. F. Huenneke, S. P. Hamburg, H. A. Mooney, *Funct. Ecol.* **2**, 335 (1988).
- P. A. T. Higgins, R. B. Jackson, J. M. Des Rosiers, C. B. Field, *Global Change Biol.* **8**, 841 (2002).
- Thanks to all of the people who helped with the JRGCE: building, harvesting, counting, and sorting. Thanks also to Jasper Ridge Biological Preserve. Supported by grants from NSF, the Morgan Family Foundation, the David and Lucile Packard Foundation, the Jasper Ridge Biological Preserve, and the Carnegie Institution of Washington. M.R.S. was supported by a Hollaender Postdoctoral Fellowship from the U.S. Department of Energy. E.S.Z. and E.E.C. received graduate support from the U.S. Environmental Protection Agency STAR Fellowship Program, the Switzer Foundation, and the A. W. Mellon Foundation. This is Carnegie Institution of Washington, Department of Global Ecology. Publication number 1.

Supporting Online Material

www.sciencemag.org/cgi/content/full/298/5600/1987/DC1
Table S1

4 September 2002; accepted 22 October 2002